

BROILER FEEDING GUIDE

Average Pounds of Feed Per Broiler Consumed During Each One Week Period

BROILER STARTER

WEEK 1	0.55 <i>lbs</i>
WEEK 2	1.00 <i>lbs</i>
WEEK 3	1.45 <i>lbs</i>
EST. TOTAL	3.00 <i>lbs each</i>

BROILER GROWER

WEEK 4	2.27 <i>lbs</i>
WEEK 5	2.80 <i>lbs</i>
WEEK 6	3.24 <i>lbs</i>
WEEK 7 <i>(optional)</i>	3.49 <i>lbs</i>
WEEK 8 <i>(optional)</i>	3.59 <i>lbs</i>
WEEK 9 <i>(optional)</i>	3.61 <i>lbs</i>
EST. TOTAL	19.00 <i>lbs each</i>



**ORGANIC
FEED MILL**

ELGIN, TEXAS

These estimates are based on test flocks raised on pasture for a full nine weeks under normal conditions. Results vary greatly, impacted by the quality of living conditions, including weather, stress, and management. Adequate access to clean water should be provided at all times. Always ensure the availability and accessibility of feed. A day without feed will result in weight loss, inhibit growth, and increase aggression; please plan your feed purchases accordingly.

For any additional information, please contact:
mill@coyotecreekfarm.org



EST. FINISH WEIGHT *Approx.* **8–10 *lbs***

DRESS WEIGHT *Approx.* **70% (of Est. Finish Weight)**



**COYOTE
CREEK
FARM**

www.CoyoteCreekFarm.org